

Predation by an Oliver's Parrot Snake *Leptophis coeruleodorsus* on a Veined Tree Frog *Trachycephalus typhonius* in Trinidad, West Indies

Oliver's Parrot Snake *Leptophis coeruleodorsus*, previously *L. ahaetulla coeruleodorsus* (Murphy *et al.* 2013), is a large, slender, arboreal and diurnal snake inhabiting forests and forest edges of eastern Colombia, Venezuela and Trinidad and Tobago (Murphy 1997, Boos 2001, Mumaw *et al.* 2015, Murphy *et al.* 2018, de Albuquerque and Fernandes 2022). It reportedly preys on frogs, lizards, snakes and small birds (Mole and Urich 1894, Mole 1924, Beebe 1946, Oliver 1948, Emsley 1977, Boos 2001, Hayes 2002, Murphy *et al.* 2013, Mumaw *et al.* 2015), but few of its prey have been identified to genus or species; Oliver (1948) identified a Boddaert's Tropical Racer *Mastigodryas boddaerti* from an unknown locality, Hayes (2002) identified three species of birds in Trinidad, Murphy *et al.* (2013) identified a *Thecadactylus* gecko in Trinidad and Mumaw *et al.* (2015) identified an Emerald-eyed Tree Frog *Boana crepitans* and a Greater Hatched-faced Tree Frog *Sphaenorhynchus lacteus* in Venezuela. Mumaw *et al.* (2015) also reported an Oliver's Parrot Snake attempting to prey on a Veined Tree Frog *Trachycephalus typhonius*, previously *Phrynohyas venulosa* and *T. venulosus* (Faivovich *et al.* 2005, Lavilla *et al.* 2010), in Venezuela until it was interrupted by a human observer. In this note, I provide the first documentation of an Oliver's Parrot Snake successfully preying on a Veined Tree Frog.

During the morning of 8 May 2009, a participant in a bird banding workshop heard the distress calls of a frog and spotted a large snake (approximately 1.5 m) with a large frog in its mouth at the William Beebe Tropical Research Station in Simla, Arima Valley, Trinidad (10°41'33.66"N, 61°17'21.88"W). The snake was wrapped around a branch about 8 m above the ground and the frog was grasped by its posterior half. As the frog struggled and emitted distress calls, its weight pulled the snake downward until it was hanging only by its tail (Fig. 1). I climbed up a masonry wall under the tree to obtain a closeup of photos within 2 m of the snake. When the snake detected my presence, it dropped down to a lower branch within 1 m of me and began to regurgitate the frog, which was covered with mucous secretions (Fig. 2). After taking a few more photos, I climbed down the wall and the snake resumed swallowing the frog. After completely swallowing the frog within an hour of our initial observation, the snake climbed higher up into the tree and vanished from our sight.

The snake was identified as an Oliver's Parrot Snake by the combination of its slender body, bright green back, copper lateral stripe, black stripe through the eye, and creamy white ventral surface (Fig. 2; Murphy 1997, Boos 2001,



Fig. 1. An Oliver's Parrot Snake dangling from a limb while grasping a Veined Tree Frog in its mouth.

Murphy *et al.* 2018). The frog was identified as a Veined Tree Frog by its paired lateral vocal sacs (Fig. 2), unique among Trinidad's tree frog species (Murphy 1997, Murphy *et al.* 2018).

Snakes of the genus *Leptophis* (taxonomy of the *L. ahaetulla* complex follows de Albuquerque and Fernandes 2022) specialize in preying on tree frogs of the family Hylidae, which comprised 83% of their diet ($n = 106$) in an early study of all taxa combined (Oliver 1948). Tree frogs comprised 90% of the diet ($n = 60$) of the Giant Parrot Snake *Leptophis ahaetulla* in northern Brazil (de Albuquerque *et al.* 2007) and 63% of the diet ($n = 16$) of the parrot snake *Leptophis marginatus* in northeastern Argentina (López *et al.* 2003). It is possible that Oliver's Parrot Snake likewise preys predominantly on tree frogs despite the scarcity of reports.

The Veined Tree Frog has toxic mucous secretions that



Fig. 2. An Oliver's Parrot Snake beginning to regurgitate a Veined Tree Frog. The bright green back, copper lateral stripe, black stripe through the eye and creamy white ventral surface characteristic of an Oliver's Parrot Snake are visible. The paired lateral vocal sacs of the Veined Tree Frog distinguish it from all other tree frog species in Trinidad.

are an antipredator strategy (Delfino *et al.* 2002, Rigolo *et al.* 2008). Although toxic skin secretions successfully foiled predation attempts by a Black-skinned Parrot Snake *Leptophis nigromarginatus* (Yeager *et al.* 2019), a Central American Indigo Snake *Drymarchon melanurus* (Leary and Razafindratsita 1998) and an Ashmead's Banded Cat-eyed Snake *Leptodeira ashmeadii* (Manzanilla *et al.* 1998), the Veined Tree Frog has been successfully preyed upon by several species of snakes, including the parrot snake *L. marginatus* (Prado 2003, de Albuquerque and Di-Bernardo 2005, Clegg 2015), Pacific Coast Parrot Snake *Leptophis diplotropis* (García-Mata *et al.* 2020), Mexican Parrot Snake *Leptophis mexicanus* (Henderson *et al.* 1977), Yellow-bellied *Liophis Erythrolamprus poecilogyrus* (da Silva *et al.* 2003), the Brown Sipo *Chironius fuscus* and potentially other species of the genus *Chironius* (Roberto and Souza 2020, Dias-Silva *et al.* 2021). It remains unknown why the toxic mucous secretions are an effective defense against

some snakes but not others.

The distress calls of frogs may also function as an antipredator strategy to startle a predator such as a bird or mammal, warn conspecifics of danger, or attract competing predators who might attempt to prey on the predator or steal the prey, facilitating the victim's escape (Hödl and Gollmann 1986, Schuett and Gillingham 1990, Hopkins and Folt 2019). In this instance and in four previous cases (Leary and Razafindratsita 1998, Prado 2003, Clegg 2015, Yeager *et al.* 2019) the distress calls of the Veined Tree Frog attracted the attention of humans. After hearing the distress calls of a Veined Tree Frog seized by a Central American Indigo Snake, Leary and Razafindratsita (1998) observed at least 17 Veined Tree Frog emerging within a 3 m radius from the crevices and cavities of a tree and orienting towards the distressed frog, presumably in response to its distress calls. These observations suggest that the distress calls of Veined Treefrogs may effectively warn conspecifics and attract secondary predators.

I thank the International Institute of Tropical Forestry for sponsoring my trip to Trinidad to attend a bird banding workshop organized by Joseph Wunderle. I also thank Renoir Auguste, Tommy Hamrick and Saifudeen Muhammad for confirming the identity of the snake and Renoir Auguste and Saifudeen Muhammad for confirming the identity of the frog based on photos posted on iNaturalist and John Murphy for reviewing the manuscript.

REFERENCES

- Beebe, W.** 1946. Field notes on the snakes of Kartabo, British Guiana, and Caripito, Venezuela. *Zoologica*, 31: 11-52.
- Boos, H. E.A.** 2001. The snakes of Trinidad and Tobago. Texas A & M University Press, College Station, Texas. 270 p.
- Clegg, J.R.** 2015. *Leptophis ahaetulla marginatus* (Southern Green Parrot Snake): Behaviour. *Herpetological Bulletin*, 131: 26-27.
- da Silva, N.J. Jr., Souza, I.F.E., Silva, W.V. and Silva, H.L.R.** 2003. Natural history notes. *Liophis poecilogyrus* (Trash-snake). Diet. *Herpetological Review*, 34: 69-70.
- de Albuquerque, N.R. and Di-Bernardo, M.** 2005. Natural history notes. *Leptophis ahaetulla marginatus* (Southern Green Parrot Snake). Diet. *Herpetological Review*, 36: 325.
- de Albuquerque, N.R. and Fernandes, D.S.** 2022. Taxonomic revision of the Parrot Snake *Leptophis ahaetulla* (Serpentes, Colubridae). *Zootaxa*, 5153:1-69.
- de Albuquerque, N.R., Galatti, U. and Di-Bernardo, M.** 2007. Diet and feeding behaviour of the Neotropical Parrot Snake (*Leptophis ahaetulla*) in northern Brazil. *Journal of Natural History*, 41: 1237-1243.
- Delfino, G., Brizzi, R., Nosi, D., and Terreni, A.** 2002. Serous cutaneous glands in New World hylid frogs: an ultrastructural study on skin poisons confirms phylogenetic

- relationships between *Osteopilus septentrionalis* and *Phrynohyas venulosa*. *Journal of Morphology*, 253: 176-186.
- Dias-Silva, F., Mattedi, C., Pontes, R.C. and Pereira, E.A.** 2021. Predation on the treefrog *Trachycephalus typhonius* (Linnaeus, 1758) by a vine snake (genus *Chironius*) in the Amazon rainforest of northern Brazil. *Herpetology Notes*, 14: 379-382.
- Emsley, M.G.** 1977. Snakes, and Trinidad and Tobago. *Bulletin of the Maryland Herpetological Society*, 13: 201-304.
- García-Mata, E.S., Cruz-Sáenz, D., Rodríguez-López, A., Ríos-Martínez, J.O., Hernández-Dávila, L.A., Lazcano, D. and Wilson L.D.** 2020. Notes on the herpetofauna of Western Mexico 23: predation by a Pacific Coast Parrot Snake (*Leptophis diplotropis*) on a Milky Treefrog (*Trachycephalus typhonius*) in the municipality Huejutla de Reyes, Hidalgo, Mexico. *Bulletin of the Chicago Herpetological Society*, 55: 101-105.
- Faivovich, J., Haddad, C.F.B., Garcia, P.C.A., Frost, D.R., Campbell, J.A. and Wheeler, W.C.** 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History*, 294: 6-228.
- Hayes, F.E.** 2002. Predation on birds by snakes in Trinidad and Tobago. *Living World, Journal of the Trinidad and Tobago Field Naturalists' Club*, 2002: 59-61.
- Henderson, R.W., Nickerson, M.A. and Hoevers, L.G.** 1977. Observations and comments on the feeding behavior of *Leptophis* (Reptilia, Serpentes, Colubridae). *Journal of Herpetology*, 11: 231-232.
- Hödl, W. and Gollmann, G.** 1986. Distress calls in Neotropical frogs. *Amphibia-Reptilia*, 7: 11-21.
- Hopkins, R. and Folt, B.** 2019. Screaming calls of *Leptodactylus savagei* (Smoky Jungle Frog) function as an alarm for conspecifics. *Journal of Herpetology*, 53: 154-157.
- Lavilla, E.O., Langone, J.A., Padial, J.M. and de Sá, R.O.** 2010. The identity of the crackling, luminescent frog of Suriname (*Rana typhonia* Linnaeus, 1758) (Amphibia, Anura). *Zootaxa*, 2671: 17-30.
- Leary, C.J. and Razafindratsita, V.R.** 1998. Attempted predation on a hylid frog, *Phrynohyas venulosa*, by an Indigo Snake, *Drymarchon corais*, and the response of conspecific frogs to distress calls. *Amphibia-Reptilia*, 19: 442-446.
- López, M.S., Giraud, A.R. and Arzamendia, V.** 2003. *Leptophis ahaetulla marginatus* (Southern Green Parrot-Snake): diet. *Herpetological Review*, 34: 68-69.
- Manzanilla, J., La Marc, E., Villareal, O. and Sanchez, D.** 1998. Natural history notes. *Phrynohyas venulose* (Veined Treefrog, "Rana Lechosa"). *Antipredator device*. *Herpetological Review*, 29: 39-40.
- Mole, R.R.** 1924. The Trinidad snakes. *Proceedings of the Zoological Society of London*, 94: 235-278.
- Mole, R.R. and Urich, E.W.** 1894. Biological notes upon some of the ophidia of Trinidad B.W.I., with a preliminary list of the species recorded from the island. *Proceedings of the Zoological Society of London*, 499-518.
- Mumaw, M.N., González, L.F.E. and Fernández, M.C.** 2015. Atlas serpientes de Venezuela: una visión actual de su diversidad. Dimacofi Negocios Avanzados S.A., Santiago, Chile. 441 p.
- Murphy, J.C.** 1997. Amphibians and reptiles of Trinidad and Tobago. Krieger Publishing Company, Malabar, Florida. 245 p.
- Murphy, J.C., Downie, J.R., Smith, J.M., Livingstone, S.M., Mohammed, R.S., Lehtinen, R.M., Eyre, M., Sewlal, J-A.N., Noriega, N., Casper, G.S., Anton, T., Rutherford, M.G., Braswell, A.L. and Jowers, M.J.** 2018. A field guide to the amphibians & reptiles of Trinidad & Tobago. Trinidad & Tobago Field Naturalists' Club, Port of Spain, Trinidad. 336 p.
- Murphy, J.C., Charles, S.P., Lehtinen, R.M., and Koeller, K.L.** 2013. A molecular and morphological characterization of Oliver's Parrot Snake, *Leptophis coeruleodorsus* (Squamata: Serpentes: Colubridae) with the description of a new species from Tobago. *Zootaxa*, 3718: 561-574.
- Oliver, J.A.** 1948. The relationships and zoogeography of the genus *Thalerophis* Oliver. *Bulletin of the American Museum of Natural History*, 92: 157-280.
- Prado, C.P.A.** 2003. Natural history notes. *Leptodactylus chaquensis* (NCN), *Pseudis paradoxa* (Paradox Frog), and *Phrynohyas venulosa* (Veined Treefrog). Predation. *Herpetological Review*, 34: 231-233.
- Rigolo, J.R., Almeida, J.A., and Ananias, F.** 2008. Histochemistry of skin glands of *Trachycephalus aff. venulosus* Laurenti, 1768 (Anura, Hylidae). *Micron*, 39: 56-60.
- Roberto, I.J. and Souza, A.R.** 2020. Review of prey items recorded for snakes of the genus *Chironius* (Squamata, Colubridae), including the first record of *Osteocephalus* as prey. *Herpetology Notes*, 13: 1-5.
- Schuett, G.W. and Gillingham, J.C.** 1990. The function of scream calling in nonsocial vertebrates: testing the predator attraction hypothesis. *Bulletin of the Chicago Herpetological Society*, 25: 137-142.
- Yeager, J., Zarlino, A., and Rodríguez, C.** 2019. Successful multimodal amphibian defence in the neotropical frog *Trachycephalus*, including handling and recovery costs to would-be predators. *Herpetology Notes*, 12: 279-280.

Floyd E. Hayes

Department of Biology, Pacific Union College, 1 Angwin Ave., Angwin, CA 94508, USA
 floyd_hayes@yahoo.com