CHECKLIST OF BIRDS OF BUSH BUSH FOREST

by C. BROOKE WORTH

In the following list observations were compiled with the help of several people, namely T.H.G. Aitken, W.G. Downs D.R. Eckelberry, R.P. ffrench, Daniel (Moon) Jury, D.W. Snow, E.S. Tikasingh. Abbreviations have been used to denote habitats in the forest, as noted below:—

S = Sky above; C = Canopy; U = Understory or low trees beyond the reach of ground nets; G = on or near the ground or water (close enough to be netted).

Tinamidae: Tinamous

Crypturellus soui	WGD, THGA. Probably resident; uncommon?
Ardeidae: Herons, etc.	differentiation
Tigrisoma lineatum	CBW. Occasional in Moriche fringe G,U.
	THGA In Petit Bush Bush Swamp
Cathartidae: New World	at end of ramp
Coragyps atratus	DWS, RPff. Common; nest found in forest.
Cathartes aura	DWS, RPff. Common; feeding in
	forest. Breeding? S,U,G.
Accipitridae: Hawks, etc	
Elanoides forficatus	DWS. Rare, one seen S
Leptodon cayanensis	DWS, Rare, one seen S
Ictinia plumbea	DWS, RPff. Fairly often seen in
	pairs; breeding? S,C.
Heterospizias meridionalis	CBW Seen once overhead S
	THGA Seen in Moriche C
Buteo nitidus	DWS, RPff. Seen occasionally. S
Leucoprernis albicollis	CBW, Rare; resident? U
Busarellus nigricollis	CBW, DRE. Rare; one pair
	seen. U
Buteogallus anthracinus	CBW, THGA. Rare; seen once. S
Spizaetus ornatus	DWS. Rare; one seen. C
Cracidae: Guans	DWS. Rare, one seen.
Pipile pipile	DMI Paras one care by same
Tipue pipue	DMJ. Rare; one seen by camp 28/3/63 and 25/4/63
Aramidae: Limpkins	TOTAL SE SING SOUTH
Aramus guarauna	DWS, CBW. A few always present; breeding? G,U.



Trinidad Piping-Guan or Pawi (Pipile pipile)




Photo by: H. DRYSDALE.

A Japanese mist net hoisted to the canopy of the forest to catch birds which seldom come to the forest floor.

THGA, CBW. Occasional in Moriche fringe. Big Bush Bush camp in dry season. Columbidae: Doves, etc. Columbigallina talpacoti nest found. Claravis pretiosa Leptotila rufaxilla CBW, RPff. Quite common; nest found. CBW, RPff. Quite common; probably breeding G,U. Psittacidae: Parrots, etc. Ara ararauna DWS, CBW. A few seen overhead. S DWS, RPff. Common; probably breeding. U Pionus menstruus Pionus menstruus DWS, RPff. Rather uncommon; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Ciccaba virgata Ciccaba virgata Ciccaba virgata Ciccaba virgata Chaetura spinicauda Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta THGA, CBW. Occasional in Morice componing of the passion. CBW, RPff. Quite common; U CBW, RPff. Quite common; probably breeding G,U. CBW, RPff. Common; probably breeding. CBW, RPff. Common; probably breeding. CDWS, RPff. Abundant resident; probably breeding. CBW, RPff. Sparse resident, probably breeding. G,U. CBW, RPff. Resident in Moriche fringe. CBW. One heard. CBW. Several seen, others often heard; probably breeding. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Several netted. CBW. Several netted. CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S CBW. One flock seen. S RPff. One seen overhead. S DWS, RPff. Rather uncommon; resident? CBV. CBW. RPff. Rather uncommon; resident? CBV. CBW. CBW. Seprendered. CBW. CBW. Seprendered. CBW. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S DWS, RPff. Rather uncommon; resident?	Jacanidae: Jacanas	and the second second
Columbidae: Doves, etc. Columbigallina talpacoti nest found. Claravis pretiosa RPff. Scen once. U Leptotila rufaxilla CBW, RPff. Quite common; nest found. U RPff. Scen once. U CBW, RPff. Quite common; probably breeding G,U. Psittacidae: Parrots, etc. Ara ararauna DWS, CBW. A few seen overhead. S Ara manilata DWS, RPff. Common; probably breeding. U Pionus menstruus DWS, RPff. Rather uncommon; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. C Cuculidae: Cuckoos, etc. Piaya minuta THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani CBW, RPff. Resident in Moriche fringe. G,U. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata THGA. One heard. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura chapmani Chaetura chapmani Chaetura chapmani Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panytila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon; CBW, RPff. Seen once. U CBW, RPff. Quite common; probably breeding G,U. CBW, RPff. Common; probably breeding. U CBW, RPff. Resident in Moriche fringe. G,U. THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. One flock seen. S RPff. One seen overhead. S DWS, RPff. Rather uncommon;	Jacana spinosa	THGA, CBW. Occasional in Moriche
Columbidae: Doves, etc. Columbigallina talpacoti nest found. Claravis pretiosa Leptotila rufaxilla Claravis pretiosa Leptotila rufaxilla Claravis pretiosa RPff. Scen once. CBW, RPff. Quite common; probably breeding G,U. Psittacidae: Parrots, etc. Ara ararauna Ara manilata DWS, CBW. A few seen overhead. S DWS, RPff. Common; probably breeding. DWS, RPff. Rather uncommon; proeding. Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Claravis pretiosa CBW, RPff. Quite common; probably breeding G,U. CBW, RPff. Quite common; probably breeding G,U. CBW, RPff. Quite common; probably breeding G,U. CBW, RPff. Common; probably breeding. CC Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident; probably breeding. CBW, RPff. Sparse resident; probably breeding. CBW, RPff. Resident in Moriche fringe. CBW, RPff. Resident in Moriche fringe. CBW, RPff. Resident in Moriche fringe. CBW. One heard. CBW. One heard. CBW. One heard. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Several seen, others often heard; probably breeding. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Seen overhead. S DWS, CBW. Several netted. CBW. One flock seen. CBW. One seen overhead. CBW. One seen overhead. CBW. One seen overhead. CBW. Several netted. CBW. One seen overhead. CBW. O		Tinge.
Claravis pretiosa Lepfotila rufaxilla Claravis pretiosa Lepfotila rufaxilla CBW, RPff. Scen once. U CBW, RPff. Quite common; probably breeding G,U. Psittacidae: Parrots, etc. Ara ararauna Ara manilata DWS, CBW. A few seen overhead. S DWS, RPff. Common; probably breeding. U DWS, RPff. Rather uncommon; breeding. C DWS, RPff. Abundant resident; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. THGA, EST. Uncommon. probably resident. G,U. CBW, RPff. Resident in Moriche fringe. G,U. CBW, RPff. Resident in Moriche fringe. G,U. THGA. Several seen, others often heard; probably breeding. U Nyctibidae: Potoos Nyctibeus griseus THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panytila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;		Big Bush Bush camp in dry season.
Claravis pretiosa Leptotila rufaxilla Claravis pretiosa Leptotila rufaxilla CBW, RPff. Quite common; probably breeding G,U. Psittacidae: Parrots, etc. Ara ararauna DWS, CBW. A few seen overhead. S DWS, RPff. Common; probably breeding. U Pionus menstruus DWS, RPff. Rather uncommon; probably breeding. C Amazona amazonica DWS, RPff. Abundant resident; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. Piaya minuta THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani Ciccaba virgata THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoproene zonaris Chaetura spinicauda Chaetura brachyura Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon; CBW. Several netted. CBW. Several netted. CBW. Several netted. CBW. Several netted. S CBW. One seen overhead. S CBW. Several netted. CBW. Se		CDW DDG Ouite common:
Claravis pretiosa Leptotila rufaxilla CBW, RPff. Quite common; probably breeding G,U. Psittacidae: Parrots, etc. Ara ararauna DWS, CBW. A few seen overhead. S DWS, RPff. Common; probably breeding. U Pionus menstruus DWS, RPff. Common; probably breeding. U Pionus menstruus DWS, RPff. Rather uncommon; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. Crotophaga ani CRW, RPff. Sparse resident, probably breeding. G,U. CBW, RPff. Resident in Moriche fringe. G,U. CBW, RPff. Resident in Moriche fringe. G,U. THGA. Several seen, others often heard; probably breeding. U Nyctibidae: Potoos Nyctibeus griseus CBW. One heard: DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Ronstantly present. S CBW. One flock seen. S RPff. One seen overhead. S DWS, RPff. Rather uncommon;	Columbigallina talpacoti	
Lepfotila rufaxilla Psittacidae: Parrots, etc. Ara ararauna DWS, CBW. A few seen overhead. DWS, RPff. Common; probably breeding. DWS, RPff. Rather uncommon; breeding. Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Rather uncommon; breeding. CBW, RPff. Sparse resident, probably breeding. CBW, RPff. Sparse resident, probably breeding. CBW, RPff. Resident in Moriche fringe. CBW. One heard. CBW. Several seen, others often heard; probably breeding. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Scen overhead. CBW. Seen overhead. CBW. Several netted. C	Claratic modicae	nest louid.
Psittacidae: Parrots, etc. Ara ararauna Namanilata Pionus menstruus Pionus menstruus Pionus menstruus Piaya cayana Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Ciccaba virgata Ciccaba virgata Apodidae: Potoos Nyctibeus griseus Porobably breeding DWS, RPff. Common; probably breeding. CBW, RPff. Rather uncommon; breeding. CBW, RPff. Sparse resident, probably breeding. CBW, RPff. Sparse resident, probably breeding. CBW, RPff. Resident in Moriche fringe. CBW, RPff. Resident in Moriche fringe. CBW, RPff. Resident in Moriche fringe. CBW. One heard. CBW. One heard. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Seen overhead. CBW. Seeral netted. CBW. Seeveral netted. CBW. Seeveral netted. CBW. Seeveral netted. CBW. Seeveral netted. CBW. One flock seen. CBW. One seen overhead. S CBW. One seen overhead. S CBW. Seeveral netted.		I(I II, Been once.
Psittacidae: Parrots, etc. Ara ararauna DWS, CBW. A few seen overhead. S DWS, RPff. Common; probably breeding. U Pionus menstruus Pionus menstruus DWS, RPff. Rather uncommon; breeding. C Amazona amazonica DWS, RPff. Abundant resident; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani CBW, RPff. Resident in Moriche fringe. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW.	серина тигахита	
Ara ararauna Ara manilata DWS, RPff. Common; probably breeding. DWS, RPff. Rather uncommon; breeding. Cuculidae: Cuckoos, etc. Piaya cayana Crotophaga ani Crotophaga ani Ciccaba virgata Ciccaba virgata CBW. A few seen overhead. DWS, RPff. Common; probably breeding. CBW, RPff. Abundant resident; breeding. CBW, RPff. Sparse resident, probably breeding. CBW, RPff. Sparse resident, probably resident. CBW, RPff. Resident in Moriche fringe. CBW. One heard. CBW. Several seen, others often heard; probably breeding. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Several netted. CBW. One flock seen. CBW. One flock seen. S RPff. Constantly present. S CBW. One flock seen. CBW. One seen overhead. CBW. Several netted. CBW. Se	Deittogidae: Parrote etc	probably breeding c,c.
Ara manilata Pionus menstruus Pionus menstruus DWS, RPff. Common; probably breeding. DWS, RPff. Rather uncommon; breeding. C DWS, RPff. Abundant resident; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Crotophaga ani Crotophaga ani Crotophaga ani Crotophaga ani Ciccaba virgata Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW, RPff. Sparse resident, probably breeding. CBW, RPff. Resident in Moriche fringe. CBW, RPff. Resident in Moriche fringe. CBW, RPff. Resident in Moriche fringe. CBW. One heard. CBW. Several seen, others often heard; probably breeding. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Several netted. CBW. One flock seen. CBW. One seen overhead. S CBW. One flock seen. CBW. One seen overhead. CBW. One seen overhead. CBW. Several netted. CBW. Several netted. CBW. One flock seen. CBW. One seen overhead. CBW. Several netted. CBW. One seen overhead. CBW. Several netted. CBW. One seen overhead. CBW. Several netted. CBW. One seen overhead. CBW. Several netted. CBW. Several neted. CBW. Several netted. CBW. Several netted. CBW. Several net		DWS CRW A few seen
Ara manilata DWS, RPff. Common; probably breeding. DWS, RPff. Rather uncommon; breeding. DWS, RPff. Abundant resident; breeding. Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Crotophaga ani Crotophaga ani Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard. CBW. One heard. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. CBW. Seen overhead. CBW. Seeral netted.	Ara arasama	· · · · · · · · · · · · · · · · · · ·
Pionus menstruus DWS, RPff. Rather uncommon; breeding. C Amazona amazonica DWS, RPff. Abundant resident; breeding. C Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani CBW, RPff. Resident in Moriche fringe. G,U. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard. U THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura chapmani Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;	Ara manilata	Overnoud.
Pionus menstruus Amazona amazonica Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Crotophaga ani Crotophaga ani Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW, CBW, CBW, CBW, CBW, CBW, CBW, CBW,	/ NI A ALIMINARA	
Amazona amazonica Dws, RPff. Abundant resident; breeding. Cuculidae: Cuckoos, etc. Piaya cayana Piaya minuta Crotophaga ani Crotophaga ani Claucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW, RPff. Sparse resident, probably breeding. G,U. THGA, EST. Uncommon. probably resident. G,U. CBW, RPff. Resident in Moriche fringe. THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Seen overhead. S Chaetura spinicauda Chaetura brachyura Chaetura cinereiventris CBW. Several netted. CBW.	Pionus menstruus	DWS. RPff. Rather uncommon;
Amazona amazonica Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. Piaya minuta THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani CBW, RPff. Resident in Moriche fringe. G,U. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura cinereiventris CBW. Several netted. CBW. Several net		
Cuculidae: Cuckoos, etc. Piaya cayana CBW, RPff. Sparse resident, probably breeding. G,U. THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani CBW, RPff. Resident in Moriche fringe. G,U. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta CBW, RPff. Sparse resident, probably resident. G,U. CBW, RPff. Resident in Moriche fringe. CBW. One heard. U THGA. One heard. U THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U THGA. Several seen, others often heard; probably breeding. U CBW. One heard; DMJ said it was vocal through dry season; probably resident. CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	Amazona amazonica	DWS, RPff. Abundant resident;
Piaya cayana Piaya minuta Piaya minuta Crotophaga ani Trotophaga ani Crotophaga ani Trotophaga ani Crotophaga ani Trotophaga ani Trot		breeding. C
Piaya minuta Piaya minuta THGA, EST. Uncommon. probably resident. G,U. CBW, RPff. Resident in Moriche fringe. G,U. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta Probably breeding. GBW. One heard. CBW. One heard; DMJ said it was vocal through dry season; probably resident. U CBW. Seen overhead. S DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S DWS, RPff. Rather uncommon;		
THGA, EST. Uncommon. probably resident. G,U. Crotophaga ani Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura cinereiventris Cypseloides niger Panyptila cayennensis THGA. Resident in Moriche fringe. THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U CBW. One heard; DMJ said it was vocal through dry season; probably resident. DWS, CBW. Seen overhead. S DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S DWS, RPff. Rather uncommon;	Piaya cayana	
Crotophaga ani Crotophaga ani		probably breeding. G,U.
Crotophaga ani CBW, RPff. Resident in Moriche fringe. Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus Nyctibeus griseus CBW. One heard, DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta CBW, RPff. Resident in Moriche fringe. G,U. THGA. One heard. U THGA. One heard, DMJ said it was vocal through dry season; probably resident. U CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	Piaya minuta	
Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U THGA. Several seen, others often heard; probably breeding. U THGA. One heard. DWS, CBW. Several probably resident. U THGA. One heard. DWS one heard. DWS seen overhead. S DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S TRACKET OF SEVERAL SERVICE S TRACKET OF SEVERAL SEVERAL S DWS, RPff. Rather uncommon;		probably resident. U.U.
Strigidae: Owls. Glaucidium brasilianum Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta THGA. One heard. U THGA. Several seen, others often heard; probably breeding. U CBW. One heard; DMJ said it was vocal through dry season; probably resident. U CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Panyptila cayennensis Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	Crotophaga ani	CBW, RPff. Resident in Moricile
Glaucidium brasilianum Ciccaba virgata THGA. One heard. THGA. Several seen, others often heard; probably breeding. Nyctibidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta THGA. One heard. U THGA. One heard. U CBW. One heard. DMJ said it was vocal through dry season; probably resident. U CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	G(2:11 O-1-	rringe.
Ciccaba virgata Nyctibiidae: Potoos Nyctibeus griseus CBW. Onc heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta THGA. Several seen, others often heard; probably breeding. U Nyctibiidae: DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts Streptoprocne zonaris DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;		TUGA One heard II
Nyctibidae: Potoos Nyctibeus griseus CBW. Onc heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta CBW. Onc heard; DMJ said it was vocal through dry season; probably resident. U CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;		1110/1. One heard.
Nyctibidae: Potoos Nyctibeus griseus CBW. One heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta CBW. One heard; DMJ said it was vocal through dry season; probably resident. U Apodidae: Swifts DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	Ciccaba virgata	
Nyctibeus griseus CBW. Onc heard; DMJ said it was vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. CBW. Onc heard; DMJ said it was vocal through dry season; probably resident. DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. Onc flock seen. S Panyptila cayennensis RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	Nyctibiidae: Potoos	neard, product, crowns.
vocal through dry season; probably resident. Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta Vocal through dry season; probably resident. U DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S Trochilidae: Hummingbirds. DWS, RPff. Rather uncommon;	Nyctibeus griseus	CBW. One heard; DMJ said it was
Apodidae: Swifts Streptoprocne zonaris Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS, CBW. Seen overhead. S DWS, RPff. Constantly present. S CBW. Several netted. CBW. Several netted. CBW. One flock seen. S RPff. One seen overhead. S TRACHICAL SERVICE SERVIC	• 6	vocal through dry season; probably
Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Panyptila cayennensis Trochilidae: Hummingbirds. Chaetura conaris CBW. Several netted. CBW. Several netted. CBW. One flock seen. SPanyptila cayennensis CBW. Several netted.		resident. U
Chaetura spinicauda Chaetura brachyura Chaetura chapmani Chaetura cinereiventris Cypseloides niger Cayennensis Trochilidae: Hummingbirds. Glaucis hirsuta DWS. Scen overhead. Scen overhead. Several netted. CBW. Several netted. CBW. One flock seen. Several netted. CBW. One seen overhead. Several netted. CBW. Several netted. CBW. One flock seen. Several netted. CBW. Several netted.		
Chaetura brachyura Chaetura chapmani Chaetura chapmani Chaetura chapmani Chaetura cinereiventris Cypseloides niger Cypseloides niger Chaetura cinereiventris CBW. Several netted. CBW. Several netted. CBW. One flock seen. S Panyptila cayennensis RPff. One seen overhead. S Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;	Streptoprocne zonaris	
Chaetura chapmani Chaetura cinereiventris Chaetura cinereiventris Cypseloides niger Chaetura cinereiventris CBW. Several netted. CBW. S		D (10. Book o termone.
Chaetura cinereiventris Cypseloides niger Cypseloides niger CBW. One flock seen. S Panyptila cayennensis RPff. One seen overhead. S Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;		
Cypseloides niger CBW. One flock seen. S Panyptila cayennensis RPff. One seen overhead. S Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;	Chaetura chapmani	
Panyptila cayennensis RPff. One seen overhead. S Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;		CRW One flock seen
Trochilidae: Hummingbirds. Glaucis hirsuta DWS, RPff. Rather uncommon;	Cypseioides niger	RPff One seen overhead S
Glaucis hirsuta DWS, RPff. Rather uncommon;	Trochilidae: Humminghi	Ki ii. Olic sceli overnead.
		DWS, RPff. Rather uncommon;
	CARRELLE STATEMENT	

Phaethornis guy	CBW, RPff. Occasionally seen;
Phaethornis longuemareus	probably resident. G,U. S DWS, RPff. Quite common;
- In the state of	probably resident. G,U.
Florisuga mellivora	DWS, CBW. Uncommon, netted.
9	Resident? G,C,U.
Chlorestes notatus	DWS, RPff. Uncommon;
•	resident? G,U.
Amazilia chionopectus	DWS, RPff. Common;
-	probably resident. G,U.
Saucerottia tobaci	DWS, RPff. Common;
	nest found. G.U.
Anthracothorax viridigula	CBW. Netted; Status unknown.
	G,U.
Anthracothorax nigricollis	
	unknown. U
Chrysolampis mosquitus	DWS, RPff. Uncommon;
\$T.\$*	resident? G,U.
Heliomaster longirostris	DWS. Doubtful sight
Trogonidae. Trogons	record. G,U.
Trogonidae: Trogons. Trogon strigilatus	CDW DDff Coorse socident
riogon strigilatus	CBW, RPff. Sparse resident, probably breeding.
Trogon violaceus	DWS, RPff. Sparse resident,
rogon violaccus	probably breeding. U
Alcedinidae: Kingfishers.	probably breeding.
Chloroceryle aenea	DWS, Fairly common in Moriche
,	fringe; netted, but status otherwise
	unknown. G.U.
Galbulidae: Jacamars.	-,
Galbula ruficauda	DWS, RPff. Relatively common;
	breeding. G,U.
Ramphastidae: Toucans	
Ramphastos vitellinus	DWS, RPff. Quite common;
Th: ' I TXY 1 1	breeding. C
Picidae: Woodpeckers	CDW DDG O
Piculus rubiginosus	CBW, RPff. Occasionally seen,
Celeus elegans	probably resident G,U. CBW. Seen once, U
Dryocopus lineatus	DWS, RPff. Sparse resident;
P E	probably breeding.
Veniliornis kirkii	DWS, CBW. Fairly common;
	probably breeding. G.U.
Phloeoceastes melanoleucos	CBW RPff. Sparse resident;
	breeding. U
Dendrocolaptidae: Woodo	

Xiphorhynchus guttatus	DWS, RPff. Common resident; breeding. G,U.
Lepidocolaptes souleyetii	CBW, RPff. Seen twice;
Deplate of the second	status unknown. G,U.
Dendrocincla fuliginosa	CBW, RPff. Common resident;
-	breeding. G,U.
Furnariidae: Ovenbirds.	
Synallaxis albescens	CBW. Seen once; status
	unknown. G
Xenops rutilans	CBW, RPff. Uncommon; probably resident. G,U.
Erundanitalan Anthindo	probably resident. G,U.
Formicariidae: Antbirds.	DWS, RPff. Very common;
Sakesphorus canadensis	breeding. G,U.
Thamnophilus doliatus	CBW, RPff. Occasionally seen;
i namnopiixus uonavas	breeding G,U.
Dysithamnus mentalis	DWS, RPff. Scattered resident;
•	breeding. G
Myrmotherula axillaris	DWS, RPff. Scattered resident;
	breeding. G
Sclateria naevia	DWS, WGD. Common near water;
	undoubtedly breeding G,U.
Myrmeciza longipes	DWS, RPff. Fairly common;
Formicarius analis	undoubtedly breeding. G THGA. Commonly heard in early
romicanus anans	morning. G
Cotingidae: Cotingas.	morning.
Attila spadiceus	DWS, RPff. Sparse resident;
-	Probably breeding. G,U.
Pachyramphus polychopte	rus CBW, RPff Occasionally
	seen. G,U.
Tityra cayana	DWS, RPff. Common resident;
This will be a distance of the same of the	probably breeding. G,U.
Pipridae: Manakins. Pipra eryhtrocephala	DWS, RPff. Uncommon; probably
т грга егуппосернага	resident. G,C,U.
Manacus manacus	DWS, WGD. A small display ground
	near 1-mile limit. G,U.
Tyrannidae: Tyrant Flyd	eatchers.
Fluvicola pica	CBW, RPff. Uncommon in
	Moriche fringe. G
Tyrannus melancholicus	DWS, RPff. Occasionally
Myjodynastas magnlatus	seen. G,U. CBW. RPff. Seen occasionally;
Myiodynastes maculatus	possibly breeding. G,U.
Megarhynchus pitangua	DWS, RPff. Seen regularly;
was brough brought	probably breeding. C,U,G.

Tyrannopsis sulphurea	DWS, RPff. Common in Moriche
	fringe; probably breeding. U,G.
Pitangus sulphuratus	CBW, RPff. Occasional in forest
	& Moriche fringe, breeding?
	Ğ,C,U.
Myiarchus tuberculifer	DWS, RPff. Present; status unknown.
Empidonax euleri	RPff. Two birds heard;
	status unknown
Platyrhinchus mystaceus	DWS, RPff. Occasionally seen;
·	probably breeding. U,G.
Tolmomyias sulphurescens	s DWS, RPff. Two records;
· -	status unknown.
Tolmomyias flaviventris	DWS, RPff. Common; probably
•	breeding. C.U.
Elaenia flavogaster	RPff. Three birds seen; status
	unknown.
Camptostoma obsoletum	DWS, RPff. Quite common;
	breeding? U,G.
Pipromorpha oleagina	CBW. One seen.
Hirundinidae: Swallows.	
Progne chalybea	DWS. Seen overhead. S
Troglodytidae: Wrens.	
Thryothorus rutilus	DWS, RPff. Common; undoubtedly
_ :	breeding. U,G.
Troglodytes aedon	DWS, RPff. Fairly common
	resident; breeding? U,G.
Turdidae: Thrushes.	-
Turdus albicollis	CBW. Rarely seen; status
G0 3	unknown. U,G.
Turdus nudigenis	CBW. Seen once. U,G.
Turdus fumigatus	CBW, RPff. Occasionally seen;
0.1."1	status unknown. II G
Sylviidae: Old World Wa	arblers.
Kamphocaenus melanurus	DWS, RPff. Occasionally seen;
\$7:	probably breeding. U,G.
Vireonidae: Vireos.	5 4446
Hylophilus aurantiifrons	DWS RPff. Common. undoubtedly
Connebide	breeding. C,U.
Coerebidae: Honeycreepe	ers.
Chlorophanes spiza	DWS, RPff. Present; status unknown.
Cyanerpes caeruleus	DWS. RPff. Occasionally seen;
Coereba flaveola	breeding? G,U.
Cocicua naveola	DWS, RPff. Common.
Parulidae: Wood Warble	Breeding. C,U,G.
Dendroica striata	
Denuroica Striata	CBW. Several netted.

Seiurus noveboracensis Geothlypis aequinoctialis	DWS Seen once. G CBW. One flew into camp house. G
Setophaga ruticilla	DWS. Seen once. G,U.
Icteridae: Blackbirds, etc	
Psarocolius decumanus	DWS, RPff. Common; breeding in
z saroconus accumunas	Moriche fringe.
Molothrus bonariensis	CBW, THGA. One nest
	parasitized. G
Icterus chrysocephalus	DWS, RPff. Fairly common in
	Moriche fringe; probably
	breeding. U,G.
Icterus nigrogularis	CBW, RPff. Fairly common in
	fringe; breeding? U,G.
Thraupidae: Tanagers.	
Euphonia violacea	DWS, RPff. Common; undoubtedly
-	breeding. G,C,U.
Tangara mexicana	CBW, RPff. Common; probably
	breeding. C,U,G.
Thraupis episcopus	RPff. Seen once. DWS, RPff. Common; undoubtedly
Thraupis palmarum	
	breeding. C,U,G.
Ramphocelus carbo	DWS, RPff. Uncommon,
_	but breeding. U,G.
Tachyphonus rufus	DWS, RPff. Uncommon, but
	probably breeding. U,G.
Tachyphonus luctuosus	CBW. Seen a few times; status unknown U,G.
The Contract of the Contract o	
Fringillidae: Finches, etc Tiaris fuliginosa	CBW. Netted; status unknown. G
Spiza americana	THGA. Many flocks flying west
Spiza americana	at dusk; one landed briefly at camp.
	27/3/63.
Sporophila intermedia	CBW. Netted; probably resident
Sporopina micrimedia	in Moriche fringe.
Sporophila lineola	CBW, RPff. Fairly common;
- K K	probably resident in Moriche
	fringe. G
Volatinia jacarina	CBW. present at N tip of
•	island. G
A total of 119 species	recorded in Bush Bush forest.

}